




# Developing satellite megaconstellation emission inventories to determine the impact on stratospheric ozone and climate.



**Impact of a decade of increasing 2019 rocket launch and re-entry emissions on stratospheric ozone depletion.**



**Contribution of individual pollutants to stratospheric  $O_3$  depletion.**




[Ryan *et al.* 2022]

**Space industry emissions cause a 0.01% decrease in global stratospheric  $O_3$ .**


**Most space industry stratospheric  $O_3$  depletion is from atmospheric re-entry  $NO_x$  (2.45 Gg in 2019).**

# Rapid increases in payload launch and re-entry rates

## Annual payloads deployed to orbit



## Annual payload re-entries



**Most payloads deployed to orbit are megaconstellation satellites.  
Short lifespan (<2 years) is already leading to increasing re-entry rates.**

## SpaceX Starlink



Launched by SpaceX Falcon 9  
Up to 60 satellites / launch  
5671 launched, 383 re-entered

## Eutelsat OneWeb



Launched by Soyuz, Falcon 9 and GSLV Mk III  
Up to 40 satellites / launch  
640 launched, 6 re-entered

**~ 540,000 extra SMC satellites planned for LEO, impacting astronomy, overcrowding and pollution.  
Environmental impacts remain under-investigated and under-regulated.**


## Launches (all atmospheric layers)



**Kerosene**  
**Falcon 9**  
LOX / RP1  
H<sub>2</sub>O  
CO  
Thermal NO<sub>x</sub>  
BC



**Hydrogen**  
**Delta IV Heavy**  
LOX / LH<sub>2</sub>  
H<sub>2</sub>O  
CO  
Thermal NO<sub>x</sub>



**Hypergolic**  
**Proton-M**  
N<sub>2</sub>O<sub>4</sub> / UDMH  
H<sub>2</sub>O  
CO  
Thermal NO<sub>x</sub>  
Fuel NO<sub>x</sub>  
BC



**Solid**  
**Long March 11**  
Al / NH<sub>4</sub>ClO<sub>4</sub> / HTPB  
H<sub>2</sub>O  
CO  
Thermal NO<sub>x</sub>  
Fuel NO<sub>x</sub>  
BC  
Chlorine  
Al<sub>2</sub>O<sub>3</sub>



**Methane**  
**Zhuque-2**  
LOX / CH<sub>4</sub>  
H<sub>2</sub>O  
CO  
Thermal NO<sub>x</sub>  
BC




## Reentries (upper atmosphere)

**Payload/Rocket**  
Thermal NO<sub>x</sub>  
Al<sub>2</sub>O<sub>3</sub>  
Other Metal Oxides?


**Most megaconstellation satellites launch using kerosene propellant.**

**Determining re-entry mass for each object is key to calculating thermal NO<sub>x</sub> and Al<sub>2</sub>O<sub>3</sub> emissions.**

**2020 - 3.27 Gg - 878 objects – 16.4% SMC**




**2021 - 4.02 Gg - 1095 objects – 20.6% SMC**



## Spatial distribution of annual object re-entries

- Megaconstellation = True, Geolocated = False
- × Megaconstellation = True, Geolocated = True
- Megaconstellation = False, Geolocated = False
- × Megaconstellation = False, Geolocated = True

**2022 - 5.58 Gg - 1650 objects – 20.9% SMC**



**Near doubling of re-entry mass since 2020 is partly driven by increasing contributions from satellite megaconstellations.**

## Reusable Objects



17.5% of re-entry mass converted to  $\text{NO}_x$ .  
No  $\text{Al}_2\text{O}_3$  emissions.

## Expendable Objects

Rocket Bodies – 70% Aluminium



Survivability:

70% Core Stage  
35% Upper Stage

Payloads – 40 % Aluminium




0% SMC Payload  
20% Non-SMC Payload

100% of re-entry mass converted to  $\text{NO}_x$ .  
 $\text{Al}_2\text{O}_3$  emissions dependent on object type.


**$\text{NO}_x$  emissions for reusable components are still based on Space Shuttle studies.  
Broad assumptions for expendable object ablation and survivability.**

# Annual emission totals for satellite megaconstellations.


## All rocket launches and re-entries




## Megaconstellation rocket launches and re-entries



## All re-entries



Similar re-entry emissions in 2021 and 2022, additional re-entries are offset by changing launch rockets. Megaconstellations contribute 9% of re-entry emissions, approaching natural injection of Al<sub>2</sub>O<sub>3</sub> and NO<sub>x</sub>.



Chemical transport model is limited by resolution and altitude (0-80km) but can monitor the impact of rocket launch / re-entry emissions on global atmospheric composition and climate.

More information is needed on emission indices and the properties of  $\text{Al}_2\text{O}_3$  aerosol from object re-entry.

- **Emission inventories for SMC and non-SMC emissions have been compiled for 2020-2022.**
  - 0.94 and 4.00 Gg of  $\text{Al}_2\text{O}_3$  and  $\text{NO}_x$  were released into the upper atmosphere in 2022.
  - Megaconstellations contribute  $\sim 9\%$  of total re-entry  $\text{Al}_2\text{O}_3$  (0.09 Gg) and  $\text{NO}_x$  (0.38 Gg) emissions in 2022.
  - Increased rocket stage reusability has mitigated the impact of increasing megaconstellation re-entries.
- **More research/data is needed to address large uncertainties:**
  - Mass of  $\text{Al}_2\text{O}_3$  /  $\text{NO}_x$  emissions from reusable re-entries.
  - % survivability and chemical composition for each re-entering object.
  - Geolocation and timestamp information for every re-entering object.
  - Increased data availability from rocket manufacturers to aid research.
  - Particle size, mass distribution and optical properties of  $\text{Al}_2\text{O}_3$  aerosol from object re-entry for modelling.
- **Next steps:**
  - Build the 2023 emission inventory.
  - Use the 2020-2022 growth rate and list of proposed constellations to predict future satellite megaconstellation emissions.
  - Simulate the impact of a decade (2020-2029) of megaconstellation emissions on stratospheric ozone and climate.

