


# Environmental Risks of Re-entry Ablation



**Eloise Marais**, [e.marais@ucl.ac.uk](mailto:e.marais@ucl.ac.uk), <https://maraisresearchgroup.co.uk/>



**Ablation layer (50-80 km)**


**Ozone layer (12-50 km)**

**10% of particles in stratosphere contaminated with space junk**

[Murphy et al., 2023]

# Greatest Concern: Slow or Reverse Ozone Recovery


## Satellite record of stratospheric ozone over the Antarctic for 1979-2018



\* no data in 1995

# Inventory of Re-entry Emissions to Track Trends

## Global re-entry emissions of pollutants from 2020 to 2024



- Nitrogen oxides
- Aluminum oxides
- Chlorine
- Black carbon

Recent increases of ~20% per year due to re-entry of megaconstellation satellites

Many metals (titanium, copper, zinc, lead, lithium, etc.) not shown

UCL spacecraft emissions tracker: <https://cbarker211.github.io/reentries.html>

Mass of some of the metals from space junk **greater than natural** input from meteors

# Combined Impacts of Re-entry Ablation and Launch Pollution

## Kerosene (SpaceX)



Pollutants injected into  
all atmospheric layers  
by rockets:

**Nitrogen oxides**

**Water vapour**

**Black carbon (BC)**

- Potent climate forcer
- Warms the atmosphere
- Warming speeds up ozone loss reactions
- Warming alters atmospheric circulation patterns

# Additional Issues and Useful Links

- Large reliance on models to gain insights into global environmental impacts
- Need observations to evidence impact, and to improve and validate models
- Observations are costly, so requires dedicated sustained funding
- Murphy et al., 2023, <https://www.pnas.org/doi/10.1073/pnas.2313374120>

**UCL Launch Emissions Tracker:**



**UCL Re-entry Emissions Tracker:**

