

Impact of UK farming on air quality, health and habitats



U. York Seminar

Eloise A Marais


16 November 2023

With ... Karn Vohra, Alok Pandey, Gongda Lu, Jamie Kelly (group members)

And ... collaborators at CEH, NOAA, ULB, AER, Environ. Canada, SUNY Albany, Rothamsted

UK NH₃ emissions overwhelming from farming


Spatial distribution of NH₃ emissions




[https://naei.beis.gov.uk/data/map-uk-das?pollutant_id=21]

UK NH₃ Emissions by activity and category

by farming activity



by management category



[UK Clean Air Strategy, 2019]

UK NH₃ emissions changes compared to other precursors




Image source: <https://www.gov.uk/government/statistics/emissions-of-air-pollutants/emissions-of-air-pollutants-in-the-uk-summary>

UK NH₃ Emissions Regulations



UNECE



8% decline relative to 2005 in
2020 to 2029

16% decline relative to 2005
from 2030

Code of Good Agricultural Practice
(COGAP) for Reducing Ammonia
Emissions (print version)

Ref: PB14506

PDF, 4.93 MB, 30 pages



Alter the inventory to meet the targets





Image source: <https://www.epa.gov/agstar/how-does-anaerobic-digestion-work>

More here: <https://www.endsreport.com/article/1846831/regulatory-capture-nfu-lobbied-defra-lower-its-global-air-quality-ambitions>


Environmental Concerns over NH_3 Emissions

Impacts health as $\text{PM}_{2.5}$ precursor


Offsets ecosystem balance via nitrogen deposition and direct exposure




NH_3 Contribution to Particulate Matter (PM) Mass



NH₃ Contribution to Particulate Matter (PM) Mass




NH_3 Contribution to Particulate Matter (PM) Mass



Instruments in space measuring NH_3 column densities

IASI: Infrared Atmospheric Sounding Interferometer



Resolution: 12 km at nadir

Swath width: 2200 km

Launch date: 2006 (2012, 2018, 2024, 2031, 2038)

Years used: 2008-2018

CrIS: Cross-track Infrared Sounder



Resolution: 14 km at nadir

Swath width: 2200 km


Launch date: 2011 (2017, 2022, 2027, 2032)

Years used: 2013-2018

Fine-scale regridding of satellite observations by oversampling

Enhance the spatial resolution relative to the native resolution of the instrument by oversampling

Oversampling Technique




Weights each IASI NH₃ pixel by area of overlap and the reported uncertainty


Oversampling code: L. Zhu,
SUSTech (Zhu et al., 2017)

Oversampling technique over London

IASI ground pixel




0.1° x 0.1° (~10 km) grid



Lose time (temporal) resolution; gain spatial resolution

Multiyear means from the IASI (morning overpass) instrument

Multiyear (2008-2018) monthly means for warmer months of the year



Column densities: number of molecules from surface of Earth to top of atmosphere

Climatological mean to be consistent with bottom-up ammonia emissions

Top-down estimate of ammonia emissions

Employ simple mass balance approach:

Convert atmospheric **column concentrations** to surface **emissions** by relating the two with a **model**

ABUNDANCES




Conversion Factor




EMISSIONS

Satellite column densities



Model Concentration-to-Emission Ratio



Satellite-derived Surface Emissions

Emission




This approach possible as NH_3 has a relatively short lifetime (2-15 hours) at or near sources


Modelled concentration-to-emissions-ratio from GEOS-Chem

GEOS-Chem

3D Atmospheric Chemistry Transport Model




UK NAEI emissions
(with temporal information)



Gas phase and heterogeneous chemistry
Transport
Dry/wet deposition

Offline assimilated
meteorology






NASA GEOS-FP for 2016

IASI-derived multiyear (2008-2018) monthly mean NH₃ emissions






Focus on **Mar-Sep** when warm temperatures and clearer conditions increase sensitivity to surface NH₃

IASI: morning
overpass



Total monthly
emissions


1 Gg = 1 kilotonne



Monthly emissions for March-September from **IASI**-derived estimates sum to **271.5 Gg**

CrIS-derived multiyear (2008-2018) monthly mean NH₃ emissions

CrIS: midday
overpass




Monthly emissions for March-September from **CrIS**-derived estimates sum to **389.6 Gg**

CrIS is 43% more than IASI. Largest difference of >a factor of 2 in September.

Satellite vs inventory NH₃ emissions: spatial distribution





Comparison of months with peak emissions according to IASI and CrIS (April and July)




Large July difference over locations dominated by dairy cattle. Inventory is 27-49% less than the satellite values.

Satellite vs inventory NH₃ emissions: seasonality

Seasonality shown as emissions in each month relative to June



Regions and annual inventory emissions



Mar-Sep emission totals in each region


1 Gg = 1 kilotonne

All reproduce spring April peak (fertilizer & manure use). Only the satellites show summer July peak (dairy cattle?).

The increase in emissions in September in CrIS is spurious.

Surface network observations corroborate top-down results

Network (points) and model (background)
surface NH_3 in Mar-Sep



Points are for DELTA instruments (blue circles)

DELTA instruments support model underestimate (**NMB = -38%**)

So do passive low-cost ALPHA instruments (yellow triangles) (**NMB = -41.5%**)




Image source:

<http://www.pollutantdeposition.ceh.ac.uk/content/ammonia-network>

GEOS-Chem underestimate in surface NH_3 driven with the NAEI corroborates results from IASI

Takehome Messages So Far

- Spring and Summer peak in NH_3 emissions
- Inventories may underestimate NH_3 emissions, as missing summer peak


What's the contribution of agricultural NH_3 to urban $\text{PM}_{2.5}$?

Test Contribution of Potentially Influential Sources

Local



City



County

National



Nearby large cities




Transport




Agriculture

Regional



Mainland Europe


Global



Desert Dust

Simulate PM_{2.5} with GEOS-Chem


3D Atmospheric Chemistry Transport Model



GEOS-Chem manual: <http://acmg.seas.harvard.edu/geos/>

Assess Validity of Model using Permanent Networks


Use total PM_{2.5} observations from the Automatic Urban and Rural Network (AURN) to assess model



WHO (2021) WHO (2005), new UK target

79% of UK exceeds updated WHO guideline


Compare annual mean surface concentrations of PM_{2.5} for 2019



Consistent spatial pattern ($R = 0.66$) and variance (slope = 1.0). Model 11% less than observations

Assess Validity of Model using Permanent Networks

Use PM_{2.5} composition measurements from UKEAP and EMEP sites to assess model





Model underpredicts observed (sulfate, nitrate, ammonium) and possibly overpredicts unobserved (dust) components. Model captures variance of components from NO_x (nitrate) and ammonia (ammonium)

Assess Validity of Model using Reference Monitors


Also evaluate model skill at reproducing observed seasonality in $\text{PM}_{2.5}$

SO_4 : sulfate; NO_3 : nitrate; NH_4 : ammonium









Enhancements in cold months and when ammonia emissions from agriculture peak due to application of synthetic fertilizer in March-April

Contribution of Sources to annual PM_{2.5} in Leicester




SO₄: sulfate; **NO₃**: nitrate; **NH₄**: ammonium
BC: black carbon; **OC**: organic carbon; **DU**: dust



Colour scale for maps of PM_{2.5} in µg m⁻³



Results for Large Cities like London and Birmingham



London: 1,600 km²
Birmingham: 270 km²
Leicester: 70 km²

Lower local than rural agricultural ammonia contribution even for large UK cities

Takehome Messages So Far


- Spring and Summer peak in NH_3 emissions
- Inventories may underestimate NH_3 emissions, as missing summer peak
- Rural NH_3 large or dominant contributor to $\text{PM}_{2.5}$ in UK cities.
- Local controls have limited efficacy at addressing $\text{PM}_{2.5}$ pollution

How effective are current measures at decreasing $\text{PM}_{2.5}$?

Emission Control Options for the UK (and EU)

Legislated emissions targets (CLE)

Emissions for present-day or PD (2019) and future (2030) for legislation (CLE)




NH_3 emissions increase, as controls insufficient to curtail increases from growth in demand

Emission Control Options for the UK (and EU)

Adoption of best best, readily available technology (**MTF**)

Emissions for present-day (**2019**) and future (**2030**) for legislation (**CLE**) vs best-available technology (**MTF**)




Projections from **ECLIPSE v6b** for all but aviation (from **IPCC**)


Best technology decreases all precursors except ammonia (NH_3) by 40-48%

NH_3 controls limited to suggested rather than enforced measures

Influence of emissions controls on $\text{PM}_{2.5}$, NH_3 , and N deposition



Influence of emission controls on PM_{2.5}




Current legislation controls cause PM_{2.5} decline of at most **2 $\mu\text{g m}^{-3}$** compared to **5 $\mu\text{g m}^{-3}$** for best technology
UK grids $> 5 \mu\text{g m}^{-3}$: 79% in the PD, 58% with legislated controls, and 36% with best technology

Relating long-term exposure to PM_{2.5} to adverse health outcomes

Available curves relating PM_{2.5} to premature mortality unconstrained at PM_{2.5} < 5 µg m⁻³

UK PM_{2.5} range is 2.5-12 µg m⁻³



PHE (2014):
Public Health England report

MacIntyre et al. (2023):
doi:10.1016/j.envint.2023.107862


GEMM:
Global Exposure Mortality Model

All curves relate adult (mostly 25+ years old) premature mortality and annual mean PM_{2.5}

Relating long-term exposure to PM_{2.5} to adverse health outcomes

Recent curves combine best of 3 well-established curves (Fusion)

Recent epidemiological study in Canada (CanCHEC) provides low-concentration constraints (eSCHIF curve)



Fusion:

Burnett et al. (2022), doi:
10.1016/j.envres.2021.112245


eSCHIF:

extended Shape Constrained
Health Impact Function (Brauer et
al., 2022 US HEI report)

Fusion addresses deficiencies in
individual curves

Relating long-term exposure to PM_{2.5} to adverse health outcomes

Hybrid curve combines Fusion and CanCHEC
Approach motivated by Weichenthal et al. (2022)



Hybrid:


eSCHIF at 2.5-9.8 $\mu\text{g m}^{-3}$ and
Fusion beyond 9.8 $\mu\text{g m}^{-3}$

Weichenthal et al. (2022) transition
between curves at 5 $\mu\text{g m}^{-3}$
requiring an artificial increase in
Fusion Relative Risks

85% of UK grids use eSCHIF in the present day; 100% in future for both scenarios. None are $< 2.5 \mu\text{g m}^{-3}$

Adult premature mortality from long-term exposure to PM_{2.5}


Values for all 184 administrative areas in the UK (115 in England, 32 in Scotland, 22 in Wales, 11 in N. Ireland)




6,751 avoided early deaths with legislated controls, double that (13,269) with best available technology
Burden of disease estimates greater than past UK-focused studies and similar to those obtained with GEMM curve

Assessing Adverse Effects on Ecosystem Health

Nitrogen critical loads
[kg N (ha sensitive habitat)⁻¹ a⁻¹]



Sensitive habitat cover
[fraction]



- 13 sensitive habitats cover 38% of UK. ~60% in Scotland.
- Use very recently revised critical loads


Critical load and sensitive habitat maps from Ed C. Rowe & N. Hina at the UK Centre for Ecology & Hydrology (**UKCEH**)

Quantify annual total nitrogen wet and dry deposition in excess of critical loads

Also assess impact of ambient NH₃ on bryophytes (NH₃ > 1 µg m⁻³)


Evaluate GEOS-Chem nitrogen wet deposition

Modelled vs observed rainwater concentrations of oxidized and reduced nitrogen



Dashed lines
bound 50%
difference

Precip-Net sites



Requires correction to monthly total
GEOS-FP precipitation
Ranges from 40-50% increase to 23-
26% decrease.
GEOS-FP annual total increases by 4%

Model 15% underestimate in nitrate may be due to low bias in NO_x emissions


Wet deposition ~60% of total deposition. Unvalidated ~40% dry deposition mostly (64%) NH_3 .

High Resolution Total Nitrogen Wet + Dry Deposition

UKCEH Concentration Based Estimated Deposition at high (5 km) spatial resolution

Open vegetation

Woodlands



GEOS-Chem too coarse to resolve deposition over sensitive habitats


GEOS-Chem also doesn't account for enhanced washout over upland areas or deposition of cloud droplets to vegetation.

GEOS-Chem total N deposition 57 Gg N less than CBED

Use CBED for present day and GEOS-Chem for response to emissions controls

Figures from Rowe et al. (2022) annual UKCEH report

Ecosystem health benefits of emission controls



Values are **total**, **mean**, and **coverage** of exceedances

Crosses show grids that fall below critical loads relative to present day

According to GEOS-Chem, more than half (60%) emitted nitrogen transported offshore

Decline in N deposition with emission controls only one-third of emissions reductions

Decline below critical loads modest. Similarly modest decrease due to past controls (2010-2019)

Exposure to harmful levels of NH₃: 73% today, 75% with legislated controls, 69% with best technology

Takehome Messages


- Spring and Summer peak in NH_3 emissions
- Inventories may underestimate NH_3 emissions, as missing summer peak
- Rural NH_3 large or dominant contributor to $\text{PM}_{2.5}$ in UK cities.
- Local controls have limited efficacy at addressing $\text{PM}_{2.5}$ pollution
- Substantial improvements to public health with emission controls, especially adoption of best available measures
- Decline in harm to sensitive habitats negligible to modest

Satellite derived emissions: JGR: Atmospheres, 2021 (doi:10.1029/2021JD035237)

Urban $\text{PM}_{2.5}$: City & Environment Interactions, 2023 (doi:10.1016/j.cacint.2023.100100)

Emission controls: GeoHealth, 2023 (doi:10.1029/2023GH000910)

Health burden of fossil-fuel derived synthetic nitrogen fertilizer




Overwhelming majority of synthetic nitrogen fertilizer from natural gas


[Vohra et al., in progress]

Health burden of fossil-fuel derived synthetic nitrogen fertilizer


Annual mean PM_{2.5} [$\mu\text{g m}^{-3}$]



PM_{2.5}-attributable early deaths



Nitrogen deposition [$\text{kg ha}^{-1} \text{a}^{-1}$]



Total attributable to PM_{2.5} is 48,625. Total attributable to all oil and gas end use activities is 3,671, so synthetic nitrogen fertilizer is ~15% of all O&G end use activities