

# Assessing the efficacy of air quality policies in the Jing-Jin-Ji region using satellite observations



UNIVERSITY OF BIRMINGHAM

Gongda Lu ([gxl642@student.bham.ac.uk](mailto:gxl642@student.bham.ac.uk)), Eloïse A Marais, Zongbo Shi

School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, UK



## 1. INTRODUCTION


- ❑ The Beijing-Tianjin-Hebei (Jing-Jin-Ji) region experiences severely degraded air quality in autumn-winter due to anthropogenic emissions from various sources and stagnant conditions that lead to buildup of pollution.
- ❑ In autumn-winter 2017-2018, strict short-term mitigation strategies were imposed to address poor air quality.
- ❑ Here we use surface observations of air pollutants from the national air quality monitoring network to assess the efficacy of these short-term pollution controls and test the skill of satellite observations in detecting changes in air quality in the Jing-Jin-Ji region.

## 2. METHODOLOGY

- ❑ Validate national air quality surface observations of PM<sub>2.5</sub> with U.S. Embassy PM<sub>2.5</sub> in Beijing.
- ❑ Estimate the effect of short-term emission controls on air pollutant concentrations by comparing autumn-winter 2016-2017 (AW2016) and autumn-winter 2017-2018 (AW2017) concentrations of fine particles (PM<sub>2.5</sub>), nitrogen dioxide (NO<sub>2</sub>), sulfur dioxide (SO<sub>2</sub>), carbon monoxide (CO), and ozone (O<sub>3</sub>).
- ❑ Compare surface and satellite observation changes in concentrations of NO<sub>2</sub>, SO<sub>2</sub> and CO. NO<sub>2</sub> and SO<sub>2</sub> are from the Ozone Monitoring Instrument (OMI) and CO is from the Measurements of Pollution in the Troposphere (MOPITT).

## 3. VALIDATION OF SURFACE OBSERVATIONS

Nongzhanguan and U.S. Embassy are 2 km apart.




**Figure 1.** Comparisons of hourly PM<sub>2.5</sub> observations at Nongzhanguan (from Sinaapp) and U.S. Embassy during AW2016 and AW2017.

- Sinaapp PM<sub>2.5</sub> at Nongzhanguan is consistent with U.S. Embassy PM<sub>2.5</sub>.
- PM<sub>2.5</sub> decreased by 44% at Nongzhanguan and 46% at U.S. Embassy.

## 5. CHANGES IN SATELLITE NO<sub>2</sub>, SO<sub>2</sub> AND CO


Satellites measure column density in the troposphere (NO<sub>2</sub>), planetary boundary layer (SO<sub>2</sub>), and throughout the total column.



**Figure 3.** Relative changes in satellite observations of NO<sub>2</sub>, SO<sub>2</sub> and CO over the Jing-Jin-Ji region and its surrounding areas. Results show (AW2017 – AW2016)/AW2016.

- NO<sub>2</sub> is stable in the Jing-Jin-Ji region and increases in the adjacent area.
- Satellite observations of SO<sub>2</sub> are too noisy to yield meaningful results.
- Slight reduction in CO is extensively observed within this domain.

## 4. PRELIMINARY RESULTS FROM SURFACE DATA



**Figure 2.** Relative changes in surface observations of PM<sub>2.5</sub>, NO<sub>2</sub>, SO<sub>2</sub>, CO and O<sub>3</sub> at 72 sites within the Jing-Jin-Ji region and 49 sites in the surrounding area. Points show (AW2017 – AW2016)/AW2016. Values inset are domain means.

- Strict emission controls were successful in reducing surface concentrations of PM<sub>2.5</sub>, NO<sub>2</sub>, SO<sub>2</sub>, and CO.
- Ozone increases with a decrease in NO<sub>x</sub>, as there is less NO available to titrate ozone.

### Data availability

National Air Quality Historical Data: <https://beijingair.sinaapp.com/>  
 U.S. Embassy PM<sub>2.5</sub> Historical Data (Beijing): [https://openaq.org/#/location/Beijing%20US%20Embassy?\\_k=lupud9](https://openaq.org/#/location/Beijing%20US%20Embassy?_k=lupud9)  
 OMI NO<sub>2</sub> Data (OMNO2d): [https://disc.gsfc.nasa.gov/datasets/OMNO2d\\_V003/summary](https://disc.gsfc.nasa.gov/datasets/OMNO2d_V003/summary)  
 OMI SO<sub>2</sub> Data (OMSO2e): [https://disc.gsfc.nasa.gov/datasets/OMSO2e\\_V003/summary](https://disc.gsfc.nasa.gov/datasets/OMSO2e_V003/summary)  
 MOPITT CO Data (V7 TIR-NIR): <https://www2.acom.ucar.edu/mopitt/products>

## 6. NEXT STEPS

- ❑ Compare coincident surface and satellite observations and also analyse satellite observations of formaldehyde (HCHO) from OMI, and ammonia (NH<sub>3</sub>) from the Atmospheric Infrared Sounder (AIRS).
- ❑ Conduct additional validation of national air quality surface observations with APHH campaign measurements.
- ❑ Use the GEOS-Chem chemical transport model to assess the contribution of interannual variability in meteorology to the changes in pollutant concentrations and interpret the results from the satellite observations.