


Influence of Oil and Gas End-Use on Summertime Particulate Matter and Ozone Pollution in the Eastern US




Karn Vohra (k.vohra@ucl.ac.uk)
with P. Achakulwisut, C. Harkins, and B. McDonald

Reliance on oil and gas in the United States


Energy consumption in the United States (1776–2019)

quadrillion British thermal units



[US EIA, 2020]

PM_{2.5} concentration ($\mu\text{g m}^{-3}$) from combustion sources in the US



[Caiazzo et al, 2013]


COVID-19 lockdowns provide an assessment of contribution of road traffic to air pollution

2020 minus 2019 (April-June)

PM_{2.5}

$\Delta = -0.4 \pm 0.2 \mu\text{g m}^{-3}$



$\mu\text{g m}^{-3}$



MDA8O₃

$\Delta = -1.4 \pm 0.6 \text{ ppb}$


ppb




... But occurrence of these lockdowns was limited to early spring.
Summertime is onset of peak ozone pollution in eastern US and
biogenic isoprene emissions in southeastern US

[He et al, 2024]

NO_x modulates the oxidative fate of isoprene



Isoprene




Isoprene
peroxy
radical

NO
49%

HO₂
27%

Isoprene SOA
precursors

Formaldehyde yield
(moles HCHO per
mole isoprene)




[Marais et al, 2012]

Here, we examine the influence of emissions from end-use activities on summertime ozone and PM_{2.5} pollution


Contribution of end-use emissions to total anthropogenic emissions

We use FIVE for on-road and off-road mobile sources and NEI 2017 for all other anthropogenic sources




Dominant end-use activities: Mobile sources for CO and NO_x, diesel vehicles for BC and volatile chemical products for NMVOCs

We use state-of-the-art 3D chemical transport model to simulate surface concentrations of atmospheric components









We run nested model with and without end-use emissions

We evaluate model output against US EPA observations



Regression slopes close to unity support use of GEOS-Chem for perturbation simulations

We evaluate model output against US EPA observations




We correct for the 3-fold model overestimate in pNO₃ and pNH₄

Contribution of end-use activities to ozone and its precursors

Large contributions to summertime mean NO_x of up to 20 ppb;



NO -to- NO_2 is small (9%) but end-use NO increases the proportion of isoprene oxidized by NO rather than by HO_2 from 42:30 to 49:27



Contribution of end-use activities to ozone and its precursors

Large contributions to summertime mean NO_x of up to 20 ppb;


NO -to- NO_2 is small (9%) but end-use NO increases the proportion of isoprene oxidized by NO rather than by HO_2 from 42:30 to 49:27



Large enhancements in formaldehyde (HCHO) from higher and more prompt HCHO yields via the NO isoprene oxidation pathway

Contribution of end-use activities to ozone and its precursors

Large contributions to summertime mean NO_x of up to 20 ppb;
 NO -to- NO_2 is small (9%) but end-use NO increases the proportion of isoprene oxidized by NO rather than by HO_2 from 42:30 to 49:27




Large enhancements in formaldehyde (HCHO) from higher and more prompt HCHO yields via the NO isoprene oxidation pathway

Ozone production in this region is limited by availability of VOCs and the enhanced HCHO yields contribute to MDA8O_3



Contribution of end-use activities to PM_{2.5} and its components

Most end-use
PM_{2.5} is
anthropogenic
OA



Contribution of end-use activities to PM_{2.5} and its components



Most end-use
PM_{2.5} is
anthropogenic
OA



Acidic pNO₃
promotes
uptake of
ammonia


Contribution of end-use activities to PM_{2.5} and its components

Most end-use
PM_{2.5} is
anthropogenic
OA





Acidic pNO₃
promotes
uptake of
ammonia

Reduction in
HO₂ isoprene
oxidation
pathway





Contribution of end-use activities to PM_{2.5} and its components

Most end-use
PM_{2.5} is
anthropogenic
OA



Acidic pNO₃
promotes
uptake of
ammonia

Reduction in
HO₂ isoprene
oxidation
pathway



Net effect
exceeds 3 µg m⁻³
mainly in cities
and northeast
coast

Influence of oil and gas end-use on eastern US summertime atmospheric composition

