

Health burden of air pollution linked to each major oil and gas lifecycle stage in the US

Karn Vohra¹, Ploy Achakulwisut², Eloise A. Marais¹, Susan Anenberg³, Gongda Lu¹, Jamie Kelly¹, Colby Francoeur^{4,5}, Colin Harkins^{4,5}, Brian McDonald⁵, Tia Scarpelli⁶


¹University College London, ²Stockholm Environment Institute, ³George Washington University, ⁴University of Colorado Boulder, ⁵NOAA Earth System Research Laboratory, ⁶Carbon Mapper

1. INTRODUCTION

- US is the world's largest producer of oil and gas. Recent (mid-2010s) increases in production and lack of enforcement of regulations targeting activities such as flaring is leading to a rapid increase in air pollution.
- Previous studies have focused either only on end use or the collective sector of oil and gas production.
- Here we use air pollutant emissions for 2017 and GEOS-Chem at high spatial resolution (25-31 km) to quantify the impact of individual stages in the lifecycle on air quality and health. Stages include upstream, midstream, downstream, and end-use activities.


2. OIL AND GAS EMISSIONS

- Air pollutant emissions from the US EPA National Emissions Inventory (NEI 2017) are aggregated for each major oil and gas lifecycle stage, Fuel-based Inventory for Vehicular Emissions (FIVE) for mobile sources.
- Oil and gas activities in the US contribute to **75%** of anthropogenic emissions for **NO_x**, **91%** for **NMVOCs** and **10%** of primary **PM_{2.5}**.



3. GEOS-CHEM MODEL: SIMULATIONS AND PERFORMANCE

- We use the 3D chemical transport model **GEOS-Chem** (v13.0.0) nested over contiguous US at 0.25°×0.3125° to simulate surface concentrations of pollutants. GEOS-Chem is run with and without emissions for each major oil and gas lifecycle stage to determine its contribution to air pollutants.
- Model evaluated against observations from the US EPA Air Quality System database and the IMPROVE program. GEOS-Chem overestimates air pollutant concentrations by **19.2%** for annual mean **PM_{2.5}** and **3.5%** for annual mean **NO₂***.




We calculate NO₂* (from thermal decomposition of NO_x-reservoir compounds) to compare to measurements from chemiluminescence instruments.


4. AIR POLLUTION FROM MAJOR OIL AND GAS LIFECYCLE STAGES IN THE US

US-wide oil and gas activities contribute to a population exposure of 2.56 µg m⁻³ (26%) of PM_{2.5}, 5.36 ppbv (68%) of NO₂, and 5.45 ppbv (11%) of maximum daily 8-hour ozone (MDA8O₃).







Upstream and midstream



Downstream




End-use




5. HEALTH BURDEN ASSESSMENT

Oil and gas activities in the US are linked to **49,500** PM_{2.5}-attributable adult premature deaths, **220,400** NO₂-attributable paediatric asthma incidences and **1,820** VOCs-attributable cancer incidences.








Upstream and midstream



Downstream



End-use



FINAL REMARKS End-use activities in the US make the largest contribution to PM_{2.5}, NO₂ and MDA8O₃, but there are large VOCs emissions (~30%) from oil and gas production. All of these are linked to adverse health outcomes underscoring the need to mitigate air pollutant emissions from the whole oil and gas lifecycle. For more information, contact Karn Vohra: k.vohra@ucl.ac.uk; X (Formerly Twitter): [@kohra_thefog](https://twitter.com/kohra_thefog)