

Modelling sources of fine particles (PM_{2.5}) in UK cities



Eloise A. Marais
Jamie M. Kelly




Jordan White
Roland J. Leigh

Particles are a mix of components that persist for days

Direct emission
of PM_{2.5}
(primary)

Emission of gas-phase
precursors
(secondary)

PM_{2.5} includes a mix of components



Black carbon primary

Sulfate

Nitrate

Ammonium

} secondary

Other inorganics


Organic aerosols primary+secondary

PM_{2.5} includes local and distant sources (long atmospheric lifetime)

Stricter World Health Organization (WHO) Guideline

(<https://apps.who.int/iris/handle/10665/345329>)


WHO Annual Air Quality Guidelines [$\mu\text{g m}^{-3}$]



Source: WHO Facebook page

Simulate $\text{PM}_{2.5}$ with the 3D Model GEOS-Chem


3D Atmospheric Chemistry Transport Model



GEOS-Chem manual: <http://acmg.seas.harvard.edu/geos/>

Simulate $\text{PM}_{2.5}$ with the 3D Model GEOS-Chem

3D Atmospheric Chemistry Transport Model



GEOS-Chem manual: <http://acmg.seas.harvard.edu/geos/>

Test Contribution of Potentially Influential Sources

Local



City



County

National



Nearby large cities




Transport




Agriculture

Regional



Mainland Europe

Global




Desert Dust


Assess Validity of Model using Reference Monitors

Use total PM_{2.5} observations from the Automatic Urban and Rural Network (AURN) to assess model

Comparison of annual mean surface concentrations of PM_{2.5} for 2019




74% of UK exceeds updated WHO guideline



Consistent spatial pattern ($R = 0.66$) and variance (slope = 1.0). Model 16% less than observations

Assess Validity of Model using Reference Monitors

Use PM_{2.5} composition measurements from UKEAP and EMEP sites to assess model




Model underpredicts observed (sulfate, nitrate, ammonium) and possibly overpredicts unobserved (dust) components. Model captures variance of components from NO_x (nitrate) and ammonia (ammonium)

Assess Validity of Model using Reference Monitors




Also evaluate model skill at reproducing observed seasonality in PM_{2.5}

SO₄: sulfate; NO₃: nitrate; NH₄: ammonium








Enhancements in cold months and when ammonia emissions from agriculture peak due to application of synthetic fertilizer in March-April


Contribution of Sources to annual PM_{2.5} in Leicester



SO₄: sulfate; **NO₃**: nitrate; **NH₄**: ammonium
BC: black carbon; **OC**: organic carbon; **DU**: dust



Contribution of Sources to PM_{2.5} seasonality in Leicester





Comparable contributions from road traffic (exhaust) and shipping emissions

Agriculture largest contributor in every month except February

Mainland Europe contribution peaks in November to April, due to long atmospheric lifetime of PM_{2.5} in winter (cold temperatures, calm conditions, low planetary boundary layer)


Corroborating Evidence from Low-Cost Sensors

Low-cost network of Zephyr® sensors distributed throughout Leicester since November 2020



Corroborating Evidence from Low-Cost Sensors

December 2020



Mean \pm std dev:


$7.2 \pm 1.0 \mu\text{g m}^{-3}$

January 2021




$9.2 \pm 1.4 \mu\text{g m}^{-3}$

February 2021




$8.7 \pm 0.8 \mu\text{g m}^{-3}$




According to low-cost sensors, local sources contribute **5-11%**. Similar to the model (**3-5%**)

Results for Large Cities like London and Birmingham



SO₄: sulfate; NO₃: nitrate;
NH₄: ammonium; BC: black carbon;
OC: organic carbon; DU: dust



London: 1,600 km²

Birmingham: 270 km²

Leicester: 70 km²

Broad applicability to other cities

Only in London is local PM_{2.5} similar to agriculture

Conclusions and Acknowledgements

- Under-regulated agricultural sector dominates PM_{2.5} year-round
- Mainland Europe makes large seasonal contribution to PM_{2.5} in November to April.
- Policies targeting local sources only likely to be effective for large cities like London
- Results reinforce the need for continued and strengthened international agreements and measures to control ammonia emissions from agriculture
- Anthropogenic dust is a large source of uncertainty due to challenges representing emissions and evaluating the model

Support provided by Leicester City Council from a Defra-funded Air Quality Grant

