

# Modelling sources of fine particles (PM<sub>2.5</sub>) in UK cities



**Eloise A. Marais**  
Jamie M. Kelly




Jordan White  
Roland J. Leigh

# Particles are a mix of components that persist for days

Direct emission  
of PM<sub>2.5</sub>  
(primary)

Emission of gas-phase  
precursors  
(secondary)

PM<sub>2.5</sub> includes a mix of components



Black carbon    **primary**

Sulfate

Nitrate

Ammonium

**secondary**

Other inorganics


Organic aerosols    **primary+secondary**

PM<sub>2.5</sub> includes local and distant sources (long atmospheric lifetime)

# Stricter World Health Organization (WHO) Guideline

(<https://apps.who.int/iris/handle/10665/345329>)

WHO Annual Air Quality Guidelines [ $\mu\text{g m}^{-3}$ ]




Source: WHO Facebook page



# Simulate $\text{PM}_{2.5}$ with the 3D Model GEOS-Chem

3D Atmospheric Chemistry Transport Model




GEOS-Chem manual: <http://acmg.seas.harvard.edu/geos/>



# Simulate $\text{PM}_{2.5}$ with the 3D Model GEOS-Chem

3D Atmospheric Chemistry Transport Model



GEOS-Chem manual: <http://acmg.seas.harvard.edu/geos/>



# Test Contribution of Potentially Influential Sources

## Local



**City**



**County**

## National



**Nearby large cities**




**Transport**




**Agriculture**

## Regional



**Mainland Europe**

## Global




**Desert Dust**


# Assess Validity of Model using Reference Monitors

Use total PM<sub>2.5</sub> observations from the Automatic Urban and Rural Network (AURN) to assess model

## Comparison of annual mean surface concentrations of PM<sub>2.5</sub> for 2019



74% of UK exceeds updated WHO guideline




Consistent spatial pattern ( $R = 0.66$ ) and variance (slope = 1.0). Model 16% less than observations



# Assess Validity of Model using Reference Monitors

Use PM<sub>2.5</sub> composition measurements from UKEAP and EMEP sites to assess model




Model underpredicts observed (sulfate, nitrate, ammonium) and possibly overpredicts unobserved (dust) components. Model captures variance of components from NO<sub>x</sub> (nitrate) and ammonia (ammonium)



# Assess Validity of Model using Reference Monitors




Also evaluate model skill at reproducing observed seasonality in PM<sub>2.5</sub>

SO<sub>4</sub>: sulfate; NO<sub>3</sub>: nitrate; NH<sub>4</sub>: ammonium








Enhancements in cold months and when ammonia emissions from agriculture peak due to application of synthetic fertilizer in March-April

# Contribution of Sources to annual PM<sub>2.5</sub> in Leicester





**SO<sub>4</sub>:** sulfate; **NO<sub>3</sub>:** nitrate; **NH<sub>4</sub>:** ammonium  
**BC:** black carbon; **OC:** organic carbon; **DU:** dust





# Contribution of Sources to PM<sub>2.5</sub> seasonality in Leicester





Comparable contributions from road traffic (exhaust) and shipping emissions

Agriculture largest contributor in every month except February

Mainland Europe contribution peaks in November to April, due to long atmospheric lifetime of PM<sub>2.5</sub> in winter (cold temperatures, calm conditions, low planetary boundary layer)

# Corroborating Evidence from Low-Cost Sensors


Low-cost network of Zephyr® sensors distributed throughout Leicester since November 2020





# Corroborating Evidence from Low-Cost Sensors


December 2020



Mean  $\pm$  std dev:


$7.2 \pm 1.0 \mu\text{g m}^{-3}$

January 2021




$9.2 \pm 1.4 \mu\text{g m}^{-3}$

February 2021




$8.7 \pm 0.8 \mu\text{g m}^{-3}$




According to low-cost sensors, local sources contribute **5-11%**. Similar to the model (**3-5%**)

# Results for Large Cities like London and Birmingham



SO<sub>4</sub>: sulfate; NO<sub>3</sub>: nitrate;  
NH<sub>4</sub>: ammonium; BC: black carbon;  
OC: organic carbon; DU: dust



London: 1,600 km<sup>2</sup>

Birmingham: 270 km<sup>2</sup>

Leicester: 70 km<sup>2</sup>

Broad applicability to other cities

Only in London is local PM<sub>2.5</sub> similar to agriculture



# Conclusions and Acknowledgements

- Under-regulated agricultural sector dominates PM<sub>2.5</sub> year-round
- Mainland Europe makes large seasonal contribution to PM<sub>2.5</sub> in November to April.
- Policies targeting local sources only likely to be effective for large cities like London
- Results reinforce the need for continued and strengthened international agreements and measures to control ammonia emissions from agriculture
- Anthropogenic dust is a large source of uncertainty due to challenges representing emissions and evaluating the model

Support provided by Leicester City Council from a Defra-funded Air Quality Grant

